

**The Effect of Mycorrhizae on Nitrogen-Fixing Bacteria Associated with
the Salt Marsh Grass *Spartina patens***

A Final Report of the 1999 Tibor T. Polgar Fellowship Program

David J. Burke
Polgar Fellow

And

Dittmar Hahn
Faculty Advisor

Dept. of Biological Sciences
Rutgers University
Newark, NJ 07102

Burke, D.J. and D. Hahn. 2000. The effect of mycorrhizae on nitrogen-fixing bacteria associated with the salt marsh grass *Spartina patens*. Section II: 28 pp. In W.C. Nieder and J.R. Waldman (Eds.). Final Reports of the Tibor T. Polgar Fellowship Program, 1999. Hudson River Foundation, New York.

ABSTRACT

Soil microbes, notably nitrogen-fixing bacteria found within the rhizosphere as well as vesicular arbuscular mycorrhizae (VAM), may play an important role in salt marsh communities by enabling plants to overcome nitrogen stress and, in so doing, structuring the plant community to favor species that form microbial symbiotic associations. We explored the effect of VAM on the community of microbial nitrogen fixers associated with the rhizosphere and soil of a common high marsh grass species, *Spartina patens*. DNA extracted from rhizosphere and soil samples was used as template for PCR amplification of *nifH*, the gene that codes for the key enzyme nitrogenase, which has a phylogenetic significance and is commonly used in the analysis of uncultured populations of nitrogen-fixing bacteria. In the presence of VAM, populations of nitrogen-fixing bacteria in the rhizosphere were found to be different from populations in bulk soil at every depth (2.5, 5.0, and 7.5 cm from the surface). In the presence of VAM, population patterns in the rhizosphere were different from treatments where VAM was excluded. Suppression of VAM altered rhizosphere populations such that they came to resemble microbial populations encountered in adjacent soil, and lowered the overall genetic heterogeneity of the N-fixing communities encountered in the rooting zone of *S. patens*.

TABLE OF CONTENTS

<u>Section of Report</u>	<u>Page Number</u>
Title Page	II-1
Abstract	II-2
Table of Contents	II-3
List of Figures/Tables	II-4
Introduction	II-5
Methods	II-7
Results	II-10
Discussion	II-20
Acknowledgements	II-24
References	II-25

LIST OF FIGURES

<u>Figure</u>	<u>Page Number</u>
1- Soil Environmental Conditions	II-11
2- Pore Water Ion Analysis	II-12
3- Effect Of Treatment On Plant Performance	II-14
4- Root Distribution In Test Plants	II-15
5- VAM Colonization Of <i>Spartina patens</i>	II-16
6- Similarity Of N-Fixing Microbial Populations	II-18
7- Comparison Of RFLP Pattern Obtained From Benomyl Pots	II-19

INTRODUCTION

Salt marshes are physically harsh environments in which facilitation among species is known to be an important mechanism influencing plant succession and tolerance of abiotic conditions (Bertness and Callaway 1994, Bertness 1992). Fertilizer amendment studies have shown that one of the most singular stresses encountered by salt marsh plants is a functional nitrogen deficiency (Chalmers 1979, Valiela and Teal 1974) brought about by soil saturation that accompanies tidal flooding (Adam 1990, Bertness 1992, Valiela and Teal 1979). Although rising tidal waters saturate marsh soil and foster N-limited conditions, these waters also import N to the marsh surface and are an important source of available N for marsh plants. The significance of the tidal N subsidy decreases with increasing distance from tidal creeks, and thus, high salt marsh usually receives less tidally imported N (Adam 1990, Bertness 1992, Chalmers 1979). This frequently results in a zone of stunted "short form" *S. alterniflora* that marks the transition between low and high marsh (Bertness 1992, Chalmers 1979). The N-budget of plants within this zone depends more on nitrogen recycled within the plant and N mineralized from soil organic matter than on tidal import (DeLaune et al. 1989, Hopkinson and Schubauer 1984, White and Howes 1994).

N-fixation is another important component of the nitrogen budget of nutrient limited, high marsh systems. N-fixation by rhizosphere bacteria may also provide as much N as tidal import (DeLaune et al. 1989, White and Howes 1994). N-fixation by diazotrophic bacteria associated with the rhizosphere, root surface and root cortex of *S. alterniflora* is fueled largely by carbon leaked from the plant root which provides the microbes with a readily available carbon source (Boyle and Patriquin 1981, Patriquin and Keddy 1978, Whiting et al. 1986). Bacterial activity for some phylogenetic groups, such as sulfate reducers, is highest for bacteria physically associated with plant root surfaces, underlining the importance of root exudation to the activity of soil micro-flora (Hines et al. 1999).

Nitrogen fixation has been found to be phosphorus limited (Hayman 1986, Mikola 1986). Diazotrophic bacteria have a high demand for P that can easily exceed the requirements of the plants with which they associate. So the bacteria plants rely upon to

provide available nitrogen are P-limited, even though the plants do not appear so limited themselves. There is evidence that the P-demand of diazotrophs can be satisfied by the activity of vesicular arbuscular mycorrhizae (VAM) associations in plant roots (Hayman, 1986, Mikola 1986). Mycorrhizae are effective at acquiring and supplying P from the soil to the host plant (Hayman 1986). VAM also transport N, usually in the form of ammonium, from the soil to the host plant (Ames et al. 1983). Although salt marshes are not considered phosphorous limited, a functional P-limitation may exist for N-fixing rhizosphere bacteria, and this may underlie the apparent N-limitation in salt marsh systems. This functional limitation may be a consequence of both high bacterial demand for P as well as the exclusion of P by iron sheaths or plaques that form on salt marsh plant roots. VAM may therefore serve a similar role to that found in legume systems, where mycorrhizae indirectly increase N-fixation by eliminating the P- limitation of the associated diazotrophic bacteria (El Ghandour et al. 1996, Hamel et al. 1991, Huang et al. 1985). Since distance from tidal flooding may increase the reliance on bacterial fixation as a N-source, high marsh plants should be more likely to host a fungal symbiont.

VAM are commonly found in many salt marshes (Cooke et al. 1993, Hoefnagels et al. 1993); however, low marsh plant species, such as *Spartina alterniflora*, are typically found without mycorrhizae possibly due to resistance to VAM infection (Hoefnagels et al. 1993, Van Duin et al. 1989). On the other hand, high marsh species such as *Spartina patens* and *Distichlis spicata* are usually encountered with VAM infection (Cooke et al. 1993). The greater competitive ability of *S. patens* over *S. alterniflora* in high marsh environments (Bertness 1991) may be due, in part, to the ability of the species to form effective VAM associations. VAM formation could improve plant performance when tidal N subsidy is low and anaerobic conditions make roots more susceptible to plaque formation. The effect of VAM on the population of P-limited diazotrophic bacteria may, therefore, be critical to successful plant activity in this region of a salt marsh. The goal of this research was to explore the interaction of vesicular arbuscular mycorrhizae with nitrogen fixing organisms associated with the roots and rhizosphere soil of a common high marsh grass species, *Spartina patens*. We used molecular techniques to determine the structural similarity between root and soil associated N-fixing microbial communities. These findings establish critical, primary

linkages between plant, fungal, and microbial aspects important in nutrient cycling and community structure in salt marsh systems.

MATERIALS AND METHOD

Sampling Site. Cores of intact plants and soil were collected from the Piermont Marsh, (41° 02' 30'' N Latitude, 73° 55' 00'' W Longitude) located along the western bank of the Hudson River approximately 18 km north of New York City. Situated within Tallman Mountain State Park and part of the National Oceanic and Atmospheric Administration's (NOAA) Hudson River National Estuarine Research Reserve (NERR), Piermont Marsh is a 500 ha tidal salt marsh that experiences a tidal range of 1 m and salinity between 0-15 ppt. The marsh is characterized by *Typha* sp. and *Phragmites australis*, but large patches of *S. patens* exist in the center of the marsh

Experimental Set-up. Soil cores (10x10x10cm) weighing approximately 1 kg fresh weight, were collected on June 7, 1999. Cores were transported back to a greenhouse at Rutgers University-Newark, fitted into plastic pots and placed in an individual pail containing 5 ppt artificial seawater (Instant Ocean® Mentor, OH, USA). Pails containing potted cores were arranged randomly on a greenhouse bench and water was added to each pail so that standing water was present 5 cm below the top of the core. This was the level of water observed in holes remaining in the field after core removal. After a 1 week acclimation period, pots were randomly exposed to one of three treatments: the fungicide benomyl (50% WP; Bonide Products, Inc. Yorkville, NY, USA) dissolved in 100 ml of tap water was applied to 8 cores at a rate of 0.1 g per kg of soil, an additional 8 cores were treated with the same benomyl solution but also received a 100 ml standard Hoagland's phosphorous solution (0.35 g KH_2PO_4 per L), and 8 cores served as controls and were treated only with tap water. Benomyl has been shown to suppress VAM colonization in previous studies (Hetrick et al. 1994) and was meant to exclude VAM from two treatments so that mycorrhizal effects could be noted on the bacterial populations. Phosphorous treatments were meant to assess microbe effects in the absence of VAM but with higher soil P levels. Benomyl-phosphorous treatments received

additional phosphorous every 2 weeks at the rate described above. Benomyl was applied only once, at the start of the study.

Soil Sampling. Soil and root samples were collected from each pot using a cork corer (1 cm diameter). During sample extraction, pH, temperature and redox potential were measured using an Oakton® pH/mv meter (Cole-Parmer Instrument Company, Vernon Hills, ILL, USA). Samples were collected from 2.5, 5.0, and 7.5 cm below the surface of each core. Each sample was divided into 3-1 g portions, placed in sterile Eppendorf tubes and kept on ice until preparation. One sample collected from each depth was preserved in paraformaldehyde and reserved for biomass estimates of phylogenetic groups by *in situ* hybridization. A second sample was frozen at -80°C and kept for bacterial population analysis, and another sample was also frozen and retained for pore-water separation and analysis.

Pore Water Chemistry. Concentrations of Cl^- and SO_4^{2-} were determined by a Dionex DX-100 ion chromatograph equipped with an IonPac AS4A-SC column using an eluent of 1.8 mM Na_2CO_3 and 1.7 mM NaHCO_3 (Hess et al., 1996; Hess et al., 1997).

Plant Characteristics. To assess plant physiological status Fv/Fm ratios, a chlorophyll fluorescence measurement of photosystem II efficiency and a reliable indicator of physiological stress, were measured on all treatments using a Hansatech FMS-2 pulse amplitude modulated fluorimeter. Plant samples were dark adapted for a minimum of 15 min prior to estimation of Fv/Fm. Measurements were made 2 weeks after treatment initiation and again 4 weeks later. After 6 weeks of growth, declines in Fv/Fm ratios for all treatments were noted and coincided with a visible decline in the greenness of plant tissue. Sampling of soil cores began at this time.

Leaf tissue was harvested from a subset (15 plants) of the soil cores based upon the determined effectiveness of fungicide treatment. Five leaves were collected from each selected plant and leaf area was measured using a Li-Cor LI3000 leaf area meter (LiCOR Instruments, Ltd., Lincoln, NE, USA). Leaf dry weight was determined after drying for 4 days at 90°C. Specific leaf area was calculated by dividing leaf dry weight by leaf area. Root density was estimated from samples used for pore water analysis. After removal of pore water, gentle washing in distilled water separated the sediment and

root material that remained. Root and soil material was then dried as indicated above and the density of both fractions determined by dividing dry weight by sample volume. Water weight of the samples, the difference between the samples original wet weight and the sum of soil and root dry weights, was used as an estimate of core saturation.

Mycorrhizae. Additional samples at each depth were collected for determination of VAM colonization. Roots for VAM estimation were stained with Trypan Blue in lactic acid using a modification of the procedure outlined by Kormanick and McGraw (1982). We used a no heat method that employed 5% KOH for de-staining. Colonization was determined using a slide mount method where 30 roots were examined for each slide. VAM colonization was determined by detecting the presence of either arbuscules or hyphal coils in sample roots. Since we were interested in the effect VAM might have on the microbial populations, samples used for population estimates were selected so that substantial differences in the extent of mycorrhizal colonization existed between the cores.

Nitrogen-fixing Microbial Populations. Rhizosphere and bulk soil samples were used to extract DNA after lysing bacteria through bead beating (Widmer et al. 1999). DNA served as template for PCR amplification. Studies on nitrogen-fixing bacteria were based on the analysis of PCR amplified structural genes for the key enzyme nitrogenase (*nifH*) which has phylogenetic significance (Aguilar et al. 1998). Several regions in the *nifH* gene code for highly conserved protein stretches and have successfully been used as target sites for PCR primers. For highly specific amplification, a nested PCR based on three primers has been described and used in molecular ecology studies (Widmer et al. 1999). The resulting amplification products were used to generate restriction fragment length polymorphism (RFLP) patterns with two restriction endonucleases (*HaeIII*, *RsaI*). After gel electrophoretic separation (7.5% polyacrylamide gels) of the different fragments, the gels were stained with silver nitrate following the procedure outlined in Bassam et al (1991). Stained gels were scanned with a Hewlett Packard ScanJet 4200C (Hewlett-Packard, Palo Alto CA, USA) and the obtained images analyzed using the GelManager 1.5 program (Biosystematica). GelManager 1.5 employs the Pearson product moment coefficient (r) to generate a band similarity matrix using information about the apparent molecular weights of the bands and band spacing, the results of which

are used to construct UPGMA (unweighed pair group method with arithmetic average) dendograms. These analyses allowed us to assess shifts in the genetic heterogeneity of microbial populations, either associated with treatment or different environmental conditions.

Statistical Analysis. Two tailed t-tests were employed to assess whether differences between treatments existed for environmental parameters. Pearson product moment coefficients calculated within the GelManager 1.5 program completed statistical analysis for comparison of microbial populations patterns. Band similarities greater than 0.7 indicate that there are no significant differences between adjacent patterns, which are then assumed to be the product of the same population. Multiple analysis of variance (MANOVA) was utilized to determine whether treatment differences existed among plants for chlorophyll fluorescence measurements.

RESULTS

Pore Water Chemistry. Mean soil pH ranged from 6.0 to 6.6 for the three treatments and no significant differences were found either between treatments or with depth (Figure 1). Soil temperature (Figure 1) for the benomyl-phosphorous cores was found to be lower than the benomyl and non-treated cores but these differences were not significant. Soil temperature remained relatively constant with depth, probably the consequence of growing plants on a greenhouse bench where solar radiation affected all sides of the core. Redox potential was found to decline (Figure 1) with depth for all treatments, the result of greater core saturation. The non-treated cores were found to have significantly lower redox potentials ($p < 0.05$) at 5.0 (-160.8 ± 38.5) and 7.5 cm (-234.0 ± 22.8) as compared to both benomyl (-50.3 ± 46.4 ; -132.8 ± 36.7) and benomyl-phosphorous cores (23.1 ± 64.9 ; -156.8 ± 36.9).

Ion analysis found no significant differences between treatments for either chloride or sulfate concentrations in pore water (Figure 2) except for chloride at the 5 cm depth where non-treated pots had lower chloride concentrations than benomyl-treated pots ($p < 0.02$). Chloride concentrations ranged from 7.6 (± 1.1) to 9.1 (± 1.0) g/l at the

Figure 1: Soil Environmental Conditions. Symbols represent the following: non-treated ■, benomyl ⊙, and benomyl phosphorous ●. VAM colonization represents the percentage of roots that contained hyphal coils indicating active mycorrhizae.



